

*Citation for published version:*

Vollaard, N & Metcalfe, R 2016, 'CrossTalk Debate: High intensity interval training does/does not have a role in risk reduction or treatment of disease: Do not write off supramaximal exercise just yet', *The Journal of Physiology*.

*Publication date:*  
2016

*Document Version*  
Peer reviewed version

[Link to publication](#)

This is the peer reviewed version of the following article: Vollaard, N & Metcalfe, R 2016, 'CrossTalk Debate: High intensity interval training does/does not have a role in risk reduction or treatment of disease: Do not write off supramaximal exercise just yet' *The Journal of Physiology*., which has been published in final form at [Link to final article using the DOI]. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

**University of Bath**

## **Alternative formats**

If you require this document in an alternative format, please contact:  
[openaccess@bath.ac.uk](mailto:openaccess@bath.ac.uk)

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## Comments on Crosstalk 26: High intensity interval training does/does not have a role in risk reduction or treatment of disease

### Do not write off supramaximal exercise just yet

Niels BJ Vollaard<sup>1</sup> and Richard S Metcalfe<sup>2</sup>

<sup>1</sup>*Department for Health, University of Bath, Bath, UK*

<sup>2</sup>*School of Sport, Ulster University, Northern Ireland, UK*

Corresponding author: [n.vollaard@bath.ac.uk](mailto:n.vollaard@bath.ac.uk)

Interestingly, this debate appears to write off supramaximal SIT as a feasible exercise intervention. Intuitively this seems reasonable, as performing the most commonly studied supramaximal SIT protocols, consisting of 4-7 repeated all-out Wingate sprints, is highly fatiguing and requires strong motivation. However, a physiological justification for the design of these protocols is lacking, and several studies have demonstrated that performing SIT protocols incorporating shorter (10-20 s (Hazell *et al.*, 2010; Metcalfe *et al.*, 2012; Zelt *et al.*, 2014)) and/or fewer sprints (2-3 (Metcalfe *et al.*, 2012; Gillen *et al.*, 2014)) does not attenuate the associated health benefits. Importantly, protocols with fewer/shorter sprints are associated with substantially lower ratings of perceived exertion (Metcalfe *et al.*, 2012), and only these protocols truly achieve the much-emphasised 'time-efficiency' of SIT and HIIT protocols.

Despite decades of research demonstrating the benefits of aerobic exercise, the uptake of, and adherence to, such exercise remains low. Therefore, it is time to consider providing alternative/adjunct exercise regimes, *alongside* current recommendations, which address the common barriers to exercise participation. There is currently no experimental evidence suggesting that supramaximal SIT is unsafe and/or poorly adhered to, and dismissing out of hand this type of exercise as unsuitable for sedentary individuals, or indeed for patient populations, may result in a missed opportunity. The best exercise intervention is one that is both effective and adhered to, and for some people this could conceivably involve time-efficient supramaximal SIT.

*Competing Interests: None Declared*

### References

Gillen JB, Percival ME, Skelly LE, Martin BJ, Tan RB, Tarnopolsky MA, Gibala MJ (2014). Three minutes of all-out intermittent exercise per week increases skeletal muscle oxidative capacity and improves cardiometabolic health. *PloS one* 9: e111489.

Hazell TJ, Macpherson RE, Gravelle BM, Lemon PW (2010). 10 or 30-s sprint interval training bouts enhance both aerobic and anaerobic performance. *Eur J Appl Physiol* 110: 153-60.

Metcalfe RS, Babraj JA, Fawcner SG, Vollaard NBJ (2012). Towards the minimal amount of exercise for improving metabolic health: beneficial effects of reduced-exertion high-intensity interval training. *Eur J Appl Physiol* 112: 2767-2775.

Zelt JG, Hankinson PB, Foster WS, Williams CB, Reynolds J, Garneys E, Tschakovsky ME, Gurd BJ (2014). Reducing the volume of sprint interval training does not diminish maximal and submaximal performance gains in healthy men. *Eur J Appl Physiol* 114: 2427-36.